

DOSING FOR IBUPROFEN (Motrin or Advil)

IBUPROFEN DOSAGE: Do not give to children under 6 months of age

| Weight | Age | Infant's Ibuprofen Concentrated Drops (50mg/1.25mL) | Children's Ibuprofen Suspension (100mg/5mL) | Children's Soft Chewable Tablets (50mg each) | Junior Strength Chewable Tablets (100mg each) | Junior Strength Ibuprofen Caplets (100 mg each) |
|--------------|------------------|---|--|---|--|--|
| | | ml | ml | Tablet | Tablet | Caplet |
| 12-17 lbs | 6- 11 mos | 1.25 mL | | | | |
| 18-23 lbs | 12- 23 mos | 1.875 mL | | | | |
| 24-35 lbs | 2-3 yrs | 2.5 mL | 5 ml | 2 | 1 | |
| 36-43 lbs | 4-5 yrs | | 7.5 ml | 3 | 1 1/2 | |
| 44-59 lbs | 6-8 yrs | | 10 ml | 4 | 2 | 2 |
| 60-70 lbs | 9- 10 yrs | | 12.5 ml | 5 | 2 ½ | 2 1/2 |
| 71-95 lbs | 11 yrs | | 15 ml | 6 | 3 | 3 |

Please Note: One dose lasts 6-8 hours. Not to be used under 6 months.

- Milliliter is abbreviated as mL; 5mL equals one teaspoon (tsp).
- Don't use household teaspoons, which can vary in size.
- Aspirin should not be used in children to treat fever or pain.